

# Quality Resource Guide

## Diagnosing & Managing the Cracked Tooth Part 2: Vertical Root Fractures

### Author Acknowledgements

**LEIF K. BAKLAND, DDS**

*Emeritus Professor*

**TORY SILVESTRIN, DDS MSD MSHPE**

*Professor, Chair and  
Advanced Program Director*

Loma Linda University School of Dentistry  
Department of Endodontics  
Loma Linda, California

Drs. Bakland and Silvestrin have no relevant financial relationships to disclose.

### Educational Objectives

Following this unit of instruction, the practitioner should be able to:

1. Differentiate between vertical root fractures and other dental fractures.
2. Identify the usual symptoms of vertical root fractures.
3. Describe methods for diagnosing vertical root fractures.
4. Explain the risk factors associated with vertical root fractures.
5. Discuss treatment options for vertical root fractures.
6. Explain the prognoses for vertical root fractures.

MetLife designates this activity for **1.5 continuing education credits** for the review of this Quality Resource Guide and successful completion of the post test.

The following commentary highlights fundamental and commonly accepted practices on the subject matter. The information is intended as a general overview and is for educational purposes only. This information does not constitute legal advice, which can only be provided by an attorney.

© 2023 MetLife Services and Solutions, LLC. All materials subject to this copyright may be photocopied for the noncommercial purpose of scientific or educational advancement.

Originally published April 2017. Updated and revised March 2020 and July 2023. Expiration date: July 2026.

The content of this Guide is subject to change as new scientific information becomes available.

**ADA CERP®** | Continuing Education  
Recognition Program

Accepted Program Provider FAGD/MAGD Credit **05/01/21 - 06/31/25**.

MetLife is an ADA CERP Recognized Provider. ADA CERP is a service of the American Dental Association to assist dental professionals in identifying quality providers of continuing dental education. ADA CERP does not approve or endorse individual courses or instructors, nor does it imply acceptance of credit hours by boards of dentistry. Concerns or complaints about a CE provider may be directed to the provider or to ADA CERP at <https://ccepr.ada.org/en/ada-cerp-recognition>.

Address comments or questions to:

DentalQuality@metlife.com - or -

MetLife Dental Continuing Education  
501 US Hwy 22  
Bridgewater, NJ 08807

**Cancellation/Refund Policy:**

Any participant who is not 100% satisfied with this course can request a full refund by contacting us.

## Introduction

Dental fractures have been categorized into: (1) crown-originating fractures (COF); (2) vertical root fractures (VRF); and (3) trauma-related fractures.<sup>1</sup> The first category was described in Part 1 of this two-part Quality Resource Guide (QRG) series (Diagnosing and Managing the Cracked Tooth Part 1: Crown-Originating Fractures 3rd Ed 2023). The focus of this QRG is vertical root fractures (VRF).

Differentiating between teeth with COF and VRF is based on the clinical history and findings, and the radiographic appearance. The two types of fractures differ in etiology, origin, locations in the tooth, and directions of fracture propagation (**Table 1**). Because of these differences, diagnosis and treatment options are different for each category, as is the prognosis.

Vertical root fractures are often not diagnosed early since symptoms frequently are lacking or mild in nature. This complicates the management of teeth with VRF. In addition, the clinical presentation can be confused with other dental conditions and can lead to misdiagnosis. The goal of this QRG is to describe clinical and radiographic aspects of VRF and discuss management of these dental problems.

## Description of VRF

Vertical root fractures (VRF) occur most commonly in restored endodontically treated teeth (**Figure 1**)<sup>2</sup> They typically originate at the root apex and progress toward the crown. But the fractures can also originate in the mid-root region at the terminus of posts and in the cervical part of the root.<sup>3</sup> Although these fractures often are asymptomatic initially, they allow ingress of bacteria into the fracture lines leading to bone resorption and periodontal pocket developments.

Excessive condensation forces during obturation of root canals and extensive post space preparations (**Figure 2**) may weaken a tooth and predispose it to VRF.<sup>4,5</sup> The etiology of VRF appears to be unrelated to thermal cycling that

causes flexure of dentin, but weakening of the dentinal walls from the root canal preparation has been suggested as a contributing factor for VRF.<sup>2,6</sup> One study, however, indicated that neither rotary nor hand files create dentinal cracks.<sup>7</sup> But another study showed that access cavity preparations had a greater negative influence on tooth strength than canal preparation.<sup>8</sup> In contrast to crown-originating fractures (COF), which usually develop in a mesial-distal direction, VRF typically occur in a buccal-lingual direction.<sup>5</sup> There is a low reported prevalence of teeth with VRF (2-5%), but when found, the management of such teeth is usually extraction or possibly root amputation in multirooted teeth.<sup>9,10,11</sup> Tamse *et al.*<sup>12</sup> found 10-20% of extracted teeth had VRF.

The clinical presentation of VRF may occur some period of time after the initiation of the fracture. The undetected fracture allows ingrowth of bacteria resulting in bony and soft tissue pathosis that often manifest as a periodontal pocket (**Figure 3**).


Vertical root fractures usually start in the apical region of a root, where endodontic files may have created micro fractures during canal preparation (**Figure 4**);<sup>13,14</sup> however, recent evidence questions the role of canal preparation.<sup>7</sup> It is not known how extensive a fracture needs to be to create symptoms. Teeth most susceptible to VRF are those with

Figure 1



A vertical root fracture (VRF) in the mesial abutment tooth of a 3-unit fixed bridge - apical origin of the fracture is evident.

Figure 2



The excessively large post preparation in tooth #4, along with the type of post used, contributed to the fracturing of the root - the radiographic lesion draping around the root is characteristic of a VRF.

Table 1 - Differences between COF and VRF

	Crown-Originating Fractures (COF)	Vertical Root Fractures (VRF)
Origin	Fractures originate in the tooth crown - usually run in a mesial-distal direction	Fractures usually originate at root apex - Often run in buccal-lingual direction.
Symptoms	Variety of symptoms from absent to severe, lancinating pain	Usually mild and often described as soreness
Radiographic appearance	Not observable unless x-ray beam runs parallel to fracture line	Not observable unless x-ray beam runs parallel fracture line, but adjacent PDL and alveolar bone may show changes
Prognosis	If diagnosed early and managed properly, teeth with COF can survive for many years	Prognosis is generally poor except in teeth where root amputation may be an option

narrow mesial-distal diameters relative to wider buccal-lingual diameters. This root configuration is found in teeth with oval, ribbon shaped and kidney shaped roots (mandibular anterior teeth, maxillary and mandibular premolars, mesial roots of mandibular molars and buccal roots of maxillary molars).<sup>9</sup>

## Symptoms

Vertical root fractures allow fluid and bacterial ingress, leading to inflammation and surrounding bone loss. A tooth with VRF often creates varying degrees of biting discomfort, swelling, tenderness to percussion and palpation, and purulent drainage through the sulcus or a gingival sinus tract.<sup>4</sup> Other symptoms that are associated with VRF include deep periodontal probing defects, periodontal-type abscess formation, periapical radiolucencies, and tooth mobility. Symptoms of VRF are variable from patient to patient, and tooth to tooth. Often the signs and symptoms are difficult to interpret and may overlap with many other dental conditions such as periodontal disease and/or failed endodontic treatment. It should be noted that the degree of pain experienced by patients with VRF is often remarkably mild, resulting in delay in seeking treatment that can lead to extensive bone loss (Figure 5).


## Diagnosis

Detection of VRF usually begins with the patient reporting symptoms such as: pain (which is generally described as “soreness”); soft tissue swelling; or presence of a sinus tract. A VRF may also be detected incidentally by observation of radiographic changes of the lamina dura and periodontal ligament space around the root and its apex. A combination of a sinus tract located near the free gingival margin, combined with a deep and somewhat narrow periodontal pocket, is an indicator of a possible VRF. Clinical examination can be aided by the use of stains and transillumination. Staining the canal space

with a dye such as methylene blue allows the dye to preferentially flow into the fracture line where it may be visualized, especially with magnification. Transillumination directed into an endodontic access cavity and down the root canal may allow visualization of VRF.<sup>4,15</sup>

The time needed for development of VRF is uncertain.<sup>16</sup> It may be years after endodontic treatment, restoration of the tooth, or post placement before evidence of a fracture is clinically detected. Although there are certain findings that are indicative of VRF (dual buccal-lingual sinus tracts, “J” shaped periapical radiolucencies along the root surface (Figure 6), and narrow deep

Figure 3




(a) An apparent periodontal pocket in the furcation area was caused by a VRF (arrow) - clinically shown in (b) - fracture appears to originate in the cervical region of the root.

Figure 4



Vertical root fractures typically originate in the apical area and progress toward the crown.


Figure 5



(a) Five years following root canal treatment of tooth #31, a lesion (arrow) was noted in the furcation - the patient had minimal symptoms and preferred to postpone any corrective treatment. (b) Two years later a large bony lesion surrounded the mesial root, yet the patient had noticed only minor soreness in the region.



Figure 6



A VRF produces an apical lesion that radiographically appears to drape around the root in a "J" shaped configuration (arrows).

Figure 7



Tooth #30 has a VRF in the mesial root that has resulted in a periodontal pocket that can easily be probed to the apical area without discomfort to the patient (a gutta-percha point is used for radiographic purposes).

periodontal pockets), no one finding is singularly definitive of VRF. The periodontal pockets that form are not as narrow as those associated with a COF and can usually be probed without using an anesthetic (Figure 7).<sup>17</sup>

Visual evidence of a fracture is the best way to establish a definitive diagnosis of VRF. This may be accomplished by re-opening the original access cavity for internal examination, or externally by raising a tissue flap (see Figure 3b). A review on the diagnosis of VRF concluded that currently there is no effective non-invasive clinical technique or radiographic markers for a definitive diagnosis of VRF.<sup>18</sup>

The difficulty in diagnosing VRF is due to several factors. For most dental problems, dentists rely on radiographic imaging; however, VRF will go undetected radiographically, unless the x-ray beam runs parallel to the fracture line. Despite not being able to routinely visualize VRF radiographically, the alveolar bone appearance around a tooth may provide clues to the presence of VRF. A loss of crestal bone continuity and/or a loss of a cortical plate may indicate an apically progressing VRF that originated in the cervical area of the tooth.<sup>19</sup>


Cone beam computed tomography (CBCT) has been used to locate and diagnose VRF.<sup>20</sup> Some studies, however, question the sensitivity and specificity of CBCT used for this purpose.<sup>21,22,23</sup> Radiopacities within a field (such gutta-percha

and posts) can create scatter and artifacts, reducing the diagnostic ability of CBCT to provide information for the presence of a fracture. Since a VRF most commonly occurs in endodontically treated teeth and may be associated with a metal post in the root, CBCT scans are not as useful for their assessment as one might anticipate. Endodontically treated teeth also have radiopaque cores, and many have radiopaque crowns, all of which add further scatter to the image.

Longitudinal fractures in teeth of 95 patients were evaluated with periapical radiographs and CBCT. The latter technique (CBCT) had 4.4 times greater odds in detecting bony defects suggestive of fractures versus periapical radiographs. Teeth with vertical root fractures were also more often associated with absent bony cortical plates, showed J shaped radiolucencies, had deeper probings (>6 mm), and were usually associated with indirect restorations. In contrast, teeth with crown-originating fractures were usually associated with direct restorations, shallower probings (<6 mm), the cortical plates were intact, and CBCTs showed angular bony defects.<sup>24</sup>

Another technique to detect vertical root fractures employs magnetic resonance imaging (MRI). A benefit of that technique is that it does not involve patient radiation. In a recent study, using Micro CT as the reference standard, it was shown that CBCT and MRI are equally reliable in the detection of vertical root fractures.<sup>25</sup>

Figure 8



(a) Tooth #14 had a VRF in the MB root - the root was resected - (b) three years post-op shows good healing (while further radiographs are not available based on patient's choice, clinically the tooth is still functioning with no reported symptoms at the date of this report). (c) Radiograph of another patient's tooth (#14) which had the MB root amputated 14 years earlier because of VRF.

In a laboratory study using MRI to evaluate induced vertical root fractures in endodontically treated teeth. The study confirmed that MRI is a valid technique for identifying the presence of vertical root fractures. The evaluators were able to consistently detect vertical root fractures as small as 26 microns.<sup>26</sup>

Misdiagnosis of teeth with VRF can complicate or delay eventual definitive treatment. Because of overlapping signs and symptoms of VRF with other dental conditions, a careful and methodical approach to diagnosis must be used to reduce errors. Early identification of VRF has many benefits, including preservation of alveolar bone, which is important for any subsequent prosthetic replacement of the fractured tooth.<sup>19</sup>

## Treatment Options

Treatment for a tooth with VRF is usually extraction, or possibly root amputation in some selected multirooted cases. Outcomes following

root amputation, especially in mandibular molars, are uncertain.<sup>27,28</sup> However, Langer<sup>29</sup> following up on his earlier original report,<sup>24</sup> noted that, in contrast to mandibular molars, resection of a single root in a maxillary molar resulted in a rather good long-term prognosis. Many clinicians have confirmed that observation. Retaining a maxillary molar for as long as possible - absent risk of continuing bone loss - appears to be a prudent treatment recommendation.<sup>30</sup>

Prior to the introduction of dental implants, the primary aim of VRF management was to repair the fracture or replace the tooth with a fixed or removable prosthesis.<sup>31</sup> Current treatment options to consider include the following:

- **Bonding the vertical fracture extraorally followed by replantation of the tooth<sup>32</sup>**

The prognosis of teeth treated by bonding of the VRF segments has been shown to initially have a high success rate (88.5% at 12 months) but it progressively drops over time (59% at

60 months).<sup>33</sup> A prospective case series of re-bonded teeth with VRF, placing mineral trioxide aggregate (MTA) into the fractures, noted clinical and radiographic success at twelve months.<sup>34</sup>


- **Root amputation**

Root amputation or hemisection in mandibular teeth with VRF have not been favorably reported in the literature,<sup>27,28</sup> but clinical experience has demonstrated some positive outcomes in maxillary multirooted molars (Figure 8).

- **Autotransplantation of teeth<sup>35</sup>**

Donor teeth for this procedure would ideally be non-functioning teeth located elsewhere in the mouth. The donor teeth must have adequate periodontal ligament support and root forms that either fit to the recipient site or the socket can be modified to receive the donor teeth (Figure 9).

Figure 9



Autotransplantation - (a) Radiograph of tooth #7 in a 50-year old male who complained of a sore and loose tooth. Root canal treatment, post and core and a crown had been completed many years earlier. (b) CBCT image shows the fractured root (arrow). (c) Clinical photo after extraction of #7. (d) Photo showing tooth #12 in a crowded alignment; the patient agreed to have it transplanted to the #7 site. (e) Radiograph taken immediately after transplantation. Root canal treatment was completed 4 weeks later. (f) Photo taken 1-month post-surgery showing soft tissue healing. (g,h) Photo and radiograph taken 3 years and 10 months after transplantation. Case Courtesy: Dr. Mitsuhiro Tsukiboshi, Aichi, Japan

- **Extraction and replacement with fixed/removable prosthesis or implant**

Extraction of a tooth with a VRF often leaves a defect in the alveolar ridge that must be augmented for the replacement prosthesis or implant to function satisfactorily (**Figure 10**).

## Prognosis

Due to constant ingress of bacteria into the fracture lines, a tooth with VRF will gradually lose support due to bone resorption. Left untreated, the condition will continue to deteriorate and can result in serious damage to the alveolar bone. Success rates for fixed prostheses or implant supported crowns surpasses that of root resections,<sup>30,36</sup> but preserving a maxillary molar by resecting the fractured root may delay the eventual tooth removal.<sup>29</sup>

## Patient Information

Due to the difficulty in diagnosing a potential VRF, it is beneficial both for the patient and the dentist to have a thorough discussion when a VRF is suspected. The discussion should include methods to be used when attempting to identify the fracture, a realistic prognosis if a VRF is present, replacement options if the tooth is non-salvageable and the importance of immediate treatment to avoid further loss of supporting tissues that can complicate tooth replacement.


The possibility of a resulting VRF should be included in informed consent discussions with patients prior to any root canal treatment. Also, the possibility that a VRF may be discovered when endodontic surgery is performed should be part of the pre-surgery discussion with the patient. Many

patients appreciate a careful and clear description of how a VRF may occur and why teeth with a VRF should not be left untreated.

## Conclusion

Vertical root fractures are often noted initially when secondary radiographic changes occur. They can lead to significant bone loss if not treated in a timely manner. Diagnosis is usually based on history (root canal treatment), radiographic appearance, and clinical findings. Although occasionally amputating the root with the fracture can retain a multirooted tooth with a VRF, often teeth with a VRF must be extracted. Today, a number of options are available for management of teeth with a VRF. All the options have better outcomes if early recognition and diagnosis of the VRF is made.

**Figure 10**





## References

- Bakland LK, Silvestrin T. Diagnosing and managing the cracked tooth – Part 1. Crown-originating fractures. *MetLife Quality Resource Guide* 2017; January.
- Abou-Rass M. Crack lines: the precursors of tooth fractures - their diagnosis and treatment. *Quintessence Int Dent Dig* 1983; 14:437-47.
- Bakland LK, Tamse A. Categorization of dental fractures. In *Vertical Root Fractures in Dentistry*. A. Tamse, I. Tsesis, E. Rosen (eds). Springer, Heidelberg. 2015, Pg. 7-28.
- Johnson WT, Leary JM. Vertical root fractures: diagnosis and treatment. *Gen Dent* 1984; 32:425-9.
- Cheung W. A review of the management of endodontically treated teeth. Post, core and the final restoration. *JADA* 2005; 136:611-9.
- Baretto MS, Moraes A, Rosa RA, Moreira CH, So MV, Bier CA. Vertical root fractures and dentin defects: effects of root canal preparation, filling, and mechanical cycling. *J Endod* 2012; 38:1135-9.
- PradeepKumar AR, Shemesh H, Archana D, Versiani MA, Sousa-Neto MD, Leoni GB, Silva-Sousa YTC, Kishen A. Root canal preparation does not induce dentinal microcracks in vivo. *J Endod* 2019;45(10):1258–1264.
- Zhang Y, Liu Y, She Y, Liang Y, Xu F, Fang C. The effect of endodontic access cavities on fracture resistance of first maxillary molar using the extended finite element method. *J Endod* 2019;45(3):316–321. .
- Gher ME Jr, Dunlap RM, Anderson MH, Kuhl LV. Clinical survey of fractured teeth. *JADA* 1987; 114:174-7.
- Bergman B, Lundquist P, Sjögren U, Sundquist G. Restorative and endodontic results after treatment with cast posts and cores. *J Prosthet Dent* 1989; 61:10-5.
- Torbjörner A, Karlsson S, Odman PA. Survival rate and failure characteristics for two post designs. *J Prosthet Dent* 1995; 73:439-44.
- Tamse A, Fuss Z, Lustig J, Kaplavi J. An evaluation of endodontically treated vertically fractured teeth. *J Endod* 1999; 25:506-8.
- Singla M, Aggarwal V, Logani A, Shah N. Comparative evaluation of rotary ProTaper, Profile, and conventional stepback technique on reduction in *Enterococcus faecalis* colony-forming units and vertical root fracture resistance of root canals. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010; 109:e105-10.
- Adorno CG, Yoshioka T, Suda H. Crack initiation on the apical root surface caused by three different nickel-titanium rotary files at different working lengths. *J Endod* 2011; 37:522-5.
- Cohen S, Blanco L, Berman L. Vertical root fractures: clinical and radiographic diagnosis. *JADA* 2003; 134:434-41.
- PradeepKumar ARP, Shemesh H, Jotbilatha S, Vijayabbarathi R, Jayalaksbmi S, Kishen A. Diagnosis of vertical root fractures in restored endodontically treated teeth: A time-dependent retrospective cohort study. *J Endod* 2016; 42:1175–1180.
- Luebke RG. Vertical crown-root fractures in posterior teeth. *Dent Clin North Am* 1984; 28:883-94.
- Berman LH, Kuttler S. Fracture necrosis: Diagnosis, prognosis assessment, and treatment recommendations. *J Endod* 2010; 36:442-446.
- Lustig JP, Tamse A, Fuss Z. Pattern of bone resorption in vertically fractured, endodontically treated teeth. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2000; 90:224-7.
- Tsesis I, Rosen E, Tamse A, Taschieri S, Kfir A. Diagnosis of vertical root fractures in endodontically treated teeth based on clinical and radiographic indices: a systematic review. *J Endod* 2010; 36:1455-8.
- Corbella AS, Del Fabro M, Tamse A, Rosen E, Tsesis I, Taschieri S. Cone beam computed tomography for the diagnosis of vertical root fractures: a systematic review of the literature and meta-analysis. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2014; 118:593-602.
- Chavda R, Mannocci F, Andiapan M, Patel S. Comparing the in vivo diagnostic accuracy of digital periapical radiography with cone-beam computed tomography for the detection of vertical root fracture. *J Endod* 2014; 40:1524-9.
- Neves FS, Freitas SQ, Campos PS, Ekestubbe A, Lofthag-Hansen S. Evaluation of cone-beam computed tomography in the diagnosis of vertical root fractures: the influence of imaging modes and root canal materials. *J Endod* 2014; 40:1530-6.
- Alaugaily I, Azim AA. CBCT patterns of bone loss and clinical predictors for the diagnosis of cracked teeth and teeth with vertical root fracture. *J Endod* 2022;48(9):1100–1106.
- Groenke BR, Idiyatullin D, Gaalaas L, Petersen A, Law A, Barsness B, Royal M, Fok A, Nixdorf DR. Sensitivity and specificity of MRI versus CBCT to detect vertical root fractures using microCT as a reference standard. *J Endod* 2023 <https://doi.org/10.1016/j.joen.2023.03.011>
- Groenke BR, Idiyatullin D, Gaalaas L, Petersen A, Chew HP, Law A, Barsness B, Royal M, Ordinola-Zapata R, Fok A, Aregawi W, Nixdorf DR. Minimal detectable width of tooth fractures using magnetic resonance imaging and method to measure. *J Endod* 2022;48(11):1414-1420
- Langer B, Stein SD, Wagenberg B. An evaluation of root resections: A ten-year study. *J Periodontol* 1981; 52:719-22.
- Zafiroopoulos G-G, Hoffmann O, Kasaj A, Willershausen B, Deli G, Tatakis DN. Mandibular molar root resection versus implant therapy: a retrospective nonrandomized study. *J Oral Implantol* 2009; 35:52-62.
- Langer B. Root resections revisited. *Int J Periodont Rest Dent* 1996; 16:200–1.
- Torabinejad M, Anderson P, Bader J, Brown LJ, Chen LH, Goodacre CJ, Kattadiyil MT, Kutsenko D, Lozada J, Patel R, Petersen F, Puterman I, White SN. Outcomes of root canal treatment and restoration, implant-supported single crowns, fixed partial dentures, and extraction without replacement: a systematic review. *J Prosthet Dent* 2007; 98:285-311.
- Pitts DL, Natkin E. Diagnosis and treatment of vertical root fractures. *J Endod* 1983; 9:338-46.
- Ozer SY, Unlu G, Deger Y. Diagnosis and treatment of endodontically treated teeth with vertical root fracture: three case reports with two-year follow-up. *J Endod* 2011; 37:97-102.
- Hayashi M, Kinomoto Y, Takeshige F, Ebisu S. Prognosis of intentional replantation of vertically fractured roots reconstructed with dentin-bonded resin. *J Endod* 2004; 30:145-8.
- Taschieri S, Tamse A, Del Fabro M, Rosano G, Tsesis I. A new surgical technique for preservation of endodontically treated teeth with coronally located vertical root fractures: a prospective case series. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010; 110:e45-52.
- Tsukiboshi M. Autotransplantation of teeth: requirements for predictable success. *Dent Traumatol* 2002; 18:157-80.
- Park S-Y, Shin S-Y, Yang S-M, Kye S-B. Factors influencing the outcome of root-resection therapy in molars: a 10-year retrospective study. *J Periodontol* 2009; 80:32-40.

## POST-TEST

Internet Users: This page is intended to assist you in fast and accurate testing when completing the “Online Exam.” We suggest reviewing the questions and then circling your answers on this page prior to completing the online exam.

(1.5 CE Credit Contact Hour) Please circle the correct answer. 70% equals passing grade.

1. **Vertical root fractures frequently occur in which direction?**
  - a. Facial-lingual
  - b. Mesial-distal
  - c. Multidirectional
  - d. Horizontal
2. **The origin of a vertical root fracture is usually from:**
  - a. Enamel toward the pulp.
  - b. The pulp toward the enamel.
  - c. The apex of the root.
  - d. The cervical area.
3. **Which of the following symptoms is not common in vertical root fractures?**
  - a. Buccal soft tissue swelling
  - b. Severe pain
  - c. Soreness to chewing
  - d. Periapical radiolucency
4. **Which of the following diagnostic procedures has only a minor role in identifying teeth with vertical root fractures?**
  - a. Cone-beam computed tomography
  - b. Percussion
  - c. Periodontal probing
  - d. Transillumination
5. **Periodontal pockets associated with a tooth having a vertical root fracture have these characteristics, EXCEPT:**
  - a. They often extend to the tooth apex
  - b. They can usually be painlessly explored
  - c. They are very narrow
  - d. They resemble pockets associated with periodontal disease
6. **Which of the following treatment procedures is not recommended for a tooth with a vertical root fracture?**
  - a. Amputation of a vertical root fractured root in multi-rooted teeth
  - b. Replacement of root canal filling
  - c. Replacement with dental implant
  - d. Bonding fractured segments
7. **Which of the following teeth with vertical root fractures have the best prognosis following root amputation?**
  - a. Maxillary incisors
  - b. Mandibular premolars
  - c. Mandibular molars
  - d. Maxillary molars
8. **Which of the following examination procedures provides the most accurate diagnosis of a vertical root fracture?**
  - a. Conventional radiographic images
  - b. CBCT images
  - c. Direct visual observation
  - d. Percussion tests
9. **Which of the following types of roots are least likely to develop a vertical root fracture?**
  - a. Premolar roots
  - b. Palatal roots of maxillary molars
  - c. Mesial roots of mandibular molars
  - d. Buccal roots of maxillary molars
10. **The prevalence of reported teeth with vertical root fractures is:**
  - a. low.
  - b. moderate.
  - c. severe.
  - d. unknown.



